

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):


- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : C07K 15/28	A1	(11) International Publication Number: WO 94/26787 (43) International Publication Date: 24 November 1994 (24.11.94)
(21) International Application Number: PCT/US94/05124 (22) International Filing Date: 4 May 1994 (04.05.94) (30) Priority Data: 08/059,323        7 May 1993 (07.05.93)        US (71) Applicant: THE BOARD OF TRUSTEES OF THE LELAND STANFORD JUNIOR UNIVERSITY [US/US]; Stanford, CA 94305 (US). (72) Inventors: [REDACTED] 1670 El Camino, #105, Menlo Park, CA 94025 (US); WEISSMAN, Irving, L.; 4147 Jefferson Avenue, Redwood City, CA 94062 (US). (74) Agents: ENAYATI, Elizabeth, F. et al.; Fenwick & West, Suite 500, Two Palo Alto Square, Palo Alto, CA 94306 (US).	(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  Published With international search report. With amended claims.  	

(54) Title: METHOD FOR GENERATING CELL TYPE SPECIFIC PHAGE ANTIBODY LIBRARIES

## (57) Abstract

A method for generating monoclonal antibodies directed against previously uncharacterized and unpurified antigens on the surface of target cells in a cell population. The method includes incubating (100) a combinatorial library of antibodies expressed on the surface of filamentous phage particles with a target cell population under conditions sufficient to bind a portion of the phage particles to the target cells. The target cells and bound phage particles are then separated (102) from the unbound phage particles, and the bound phage particles are recovered (104). These phage particles are then amplified (106) to create an enriched library. Monoclonal antibodies specific to the target cell are then isolated (108) from the enriched library for subsequent use.



**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

**Method for Generating Cell Type Specific Phage Antibody Libraries**

5

**BACKGROUND OF THE INVENTION**

10 The invention was supported by a grant of the National Institute of Health/National Cancer Institute, Grant No. CA-42551, under which the government may have rights.

**Field of the Invention**

15 The invention relates generally to the field of cell type specific phage antibody libraries, and relates specifically to the field of generating such libraries by direct affinity selection on the surface of target cells.

Although the generation of antibody libraries, and more specifically phage antibody libraries, has been done for selected, purified antigens, a method for generating cell type specific phage antibody libraries for identifying, characterizing, and cloning new surface molecules on intact, whole cells has not  
20 been developed. Such cell surface molecules are likely to be mediators of important biological functions and may be used in ~~clinical applications as targets for pharmacological intervention and surface markers for identification, purification, and phenotypic dissection of the target cells.~~

**Brief Description of Background Art**

25 The phage antibody technology is a recently developed technique that allows the expression of monoclonal antibodies on the surface of filamentous phage particles (see, for example, Smith, Science 228, 1315-1317 (1985); Barbas et al., Proc. Natl. Acad. Sci. USA 88, 7978-7982 (1991); Clackson et al., Nature 352, 624-628 (1991)).

30 One advantage of this technology is the linkage of the specificity of a monoclonal antibody with the genetic information encoding the specificity in the same phage particle, thus allowing enrichment of specific phage antibodies by binding to purified antigens attached to a solid phase, followed by elution and amplification of the phage particles. Such a method is described in Kang et al.,  
35 Proc. Natl. Acad. Sci. USA 88, 4363-4366 (1991). Using ~~micropanning or affinity column techniques~~, researchers have used this technology to generate antibodies reactive to well-characterized antigens that can be purified in large quantities,

either as natural or recombinant products, including various haptens, tetanus toxoid, HIV and RSV glycoproteins, HBV core and surface antigens, progesterone, thyroglobulin, TNF alpha, CEA, soluble CD4, IL2R p55 chain, lysozyme, bovine serum albumin, trypsin, anhydrotrypsin, and ~~B6 tumor associated antigen.~~

5           However, none of these methods involve using the phage antibody technology to generate antibodies directed against previously unknown and unpurified antigens, including cell surface antigens. Accordingly, there remains a need for a method for rapidly generating monoclonal antibodies directed against unpurified surface antigens on phenotypically defined cell populations using this  
10           technology.

### SUMMARY OF THE INVENTION

          The present invention involves methods for using the phage antibody technology to generate antibodies directed against previously unknown and unpurified antigens on the cell surface, e.g., as a way to discover new antigens on  
15           important cell types. Since a phage antibody may be amplified by growing the phage in bacteria, it is possible to recover a single phage particle specifically bound to an antigen in order to generate enough monoclonal antibodies for detailed characterization of the antigen. The high sensitivity of the method enables isolation of antibodies directed against an unpurified antigen in the midst  
20           of numerous background antigens.

          In the present invention, target antigens on the surface of a phenotypically defined population of cells are used to separate phage antibodies reactive to the antigens from the background. The cell membrane of the target cells can be considered a solid phase support for a set of cell type specific antigens, and the  
25           intact target cells can be used as a convenient physical device to separate phage antibodies bound on the surface of the target cells from the unbound phage antibodies.

          The inventive method for generating antibodies against previously uncharacterized and unpurified antigens on the surface of target cells in a cell  
30           population includes incubating a combinatorial library of antibodies expressed on the surface of filamentous phage particles with a target cell population under conditions sufficient to bind a portion of the phage particles to the target cells. The target cells and bound phage particles are then separated from the unbound phage particles, and the bound phage particles are recovered. These phage  
35           particles are then amplified to create an enriched library. Monoclonal antibodies specific to the target cell are then isolated from the enriched library for subsequent use.

To selectively enrich phage antibodies directed against antigens unique to the target cells, the phage antibody library may be ~~preadsorbed~~ with an excess of non-target cells to immunologically remove antibodies directed against antigens present on ~~the target cells~~. The adsorption step may be performed after the  
5 phage antibody selection and amplification, or prior to the initial incubating step. ~~Alternatively~~, the removal of these antibodies can be done by ~~subtraction~~ hybridization of DNA from a phage antibody library enriched with the target cells to DNA from a library enriched with non-target cells. The phenotypically defined target cell population may be isolated prior to incubating the target cells  
10 with the phage antibody library, or the target cells may be a subpopulation within a heterogeneous cell population and be isolated after incubating with the phage antibody library, thus copurifying the phage antibodies bound to the cell surface antigens.

### BRIEF DESCRIPTION OF THE FIGURES

15 FIGURE 1 is a flow chart illustrating an embodiment of the present inventive method.

FIGURE 2 is a flow chart illustrating another embodiment of the present inventive method.

20 FIGURES 3a - 3d are histograms showing indirect immunofluorescence patterns of KG1a cells and mouse thymocytes stained with the unenriched and enriched antibody libraries. The abscissa indicates relative fluorescence intensity, and the ordinate indicates relative cell number.

### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

The process of the present invention includes a method to generate  
25 antibodies against cell surface antigens by direct affinity selection of phage antibodies on the surface of target cells. Generally, a phage antibody library is constructed by standard methods and an aliquot of the library is used for incubation with the target cells. Phage antibodies that are specific to and have a high affinity for antigens on the target cells will tightly bind to the cells and can be  
30 separated from the rest of the phage antibody library by washing and spinning down the cells. The bound phage particles can then be eluted from the cell surface, recovered, and amplified to yield an enriched phage antibody library.

As illustrated in the flow chart of FIGURE 1, the present method involves a first step of incubating (100) a standard phage antibody library, such as a  
35 combinatorial library of antibodies expressed on the surface of filamentous phage

particles, with a target cell population. The cells are incubated under conditions sufficient to bind a portion of the phage particles to the target cells. In the embodiment of FIGURE 1, the target cell population is an isolated cell population. The process of the present invention preferably utilizes a standard combinatorial  
5 phage antibody library of  $10^7$  recombinants, which may be prepared using standard protocols, such as that described in Barbas et al., *Methods* 2, 119-124 (1991). The cell population may be isolated according to any preselected criteria.

Following incubation (100), the ~~target cells and the phage particles bound thereto are separated (102) from the unbound phage particles.~~ Phage antibodies  
10 that are specific to and have a high affinity for antigens on the target cells will tightly bind to the cells and can be separated from the rest of the phage antibody library by centrifugating the cells through a calf serum cushion. Other standard separation methods known by those skilled in the relevant art may be used, examples including but not limited to, cell washing and centrifugation, filtration  
15 of the unbound phages through a membrane, and immobilization or capture of the cells on a solid phase. The separated phage particles, or phage antibodies, are then recovered (104). The bound phage particles may be eluted, for example using an acidic buffer, from the cell surface using standard methods. The resulting recovered phage antibodies are amplified (106) to yield an enriched  
20 phage antibody library. ~~Repeating this enrichment step several times generates a~~ phage antibody library with high specificity for surface antigens on the target cells. Each round of enrichment increases the representation of antibodies with higher affinity for the target antigens but decreases the diversity of the library. Thus, the amount of enrichment will depend on the particular requirements of the  
25 user.

Once the enriched library is generated, monoclonal antibodies specific for the target cells are isolated (108) from the library. Thus, phage antibodies reactive to antigens on the target cell population are obtained. The monoclonal antibodies are isolated (108) using methods known and available to those skilled  
30 in the art.

In one form of the invention, direct affinity selection of the phage antibodies on the cell surface is used as described substantially above. In that instance, a population of isolated target cells are incubated with the phage antibody library. The target cells may be in the form of a cell line, or other  
35 substantially isolated population of target cells.

In an alternative embodiment, and as illustrated in FIGURE 2, the target cell population is a subpopulation from a heterogeneous cell population before

any isolation. After incubation with the phage antibody library, ~~the target cell population may subsequently be separated (110) from other cells in the~~ heterogeneous population based on phenotypic distinctions, or other appropriate distinctions, thus copurifying the phage particles bound to the cell surface (112).

5            Preferably, using the copurification (110) method illustrated in FIGURE 2, the target cells are phenotypically defined target cell subpopulations. This allows generation of antibodies against rare but phenotypically defined cell subpopulations that are too few in cell number to allow characterization by conventional methods. In this embodiment of the inventive method, the phage  
10           antibody library is ~~first allowed to bind to the total and unpurified cell population.~~ The target subpopulation is then physically separated from the rest of the cells by means of its defining phenotypic characteristics. The phage particles specifically bound to the surface antigens of the target cells are separated (112) out together with the target cell subpopulation.

15           The separated phage particles can then be recovered (114), amplified (116), and used in another round of enrichment in a manner similar to that described above. Thus, phage antibodies reactive to antigens on the target subpopulation of a heterogeneous population may be selectively enriched by copurification of the target cells with their specific phage antibodies, exploiting the phenotypic  
20           characteristics of the target cells. Standard methods of cell separation known to those skilled in the art may be used.

             The method to be used for the copurification (112) step may be any of the type known and available to those skilled in the art. Exemplary methods applicable to target cells in suspension include fluorescence-activated cell sorting  
25           (FACS), immunological ~~pairing~~, magnetic bead separation, and affinity chromatography. The specific copurification method will depend on availability of materials, the type of target cell subpopulation, and other factors determinable by those skilled in the art.

             The cell population used in conjunction with the present method may  
30           include whole tissue sections, such as frozen tissue sections, in which the target cell population is identifiable. Exemplary methods for identifying and isolating target cell subpopulations may include use of immunohistochemical markers, physical location of cell populations, or other methods known and used by those skilled in the art. In practicing the present invention using tissue sections, the  
35           tissue section is incubated with the phage antibody library under conditions sufficient to bind phage particles to the target cells. The target cells and the bound phage particles are then microdissected from the tissue section, the phage



particles are recovered and amplified, and monoclonal antibodies are isolated in accordance with the method described above.

The preferred embodiment of the invention makes use of the phenotypic characteristics of the target cells to identify phage antibodies of interest by means of their binding to the target cells. In an alternative embodiment, phage antibodies are selected if they are shown to have an identifiable and preselected function, as compared with other phage antibodies without such function.

In that embodiment, phage antibodies may be assayed, for example, for the functional consequences of a phage antibody-target cell binding. One assay may be to measure intracellular calcium concentration. Another effective assay may be used to detect cell activation, cell proliferation, or cell killing, or the blocking of these activities. Yet another assay may be to analyze the expression of preselected genes. Other functional methods of identifying phage antibodies known and available to those skilled in the art may be used.

In practicing the present invention, the resulting phage antibody library may contain antibodies reactive to antigens unique to the target cells or ~~antibodies reactive to antigens present on both the target cells and non-target cells. It may be desirable to selectively remove the antibodies reactive to antigens not unique to the target cells.~~ This may be accomplished by preadsorbing the phage antibody library with an excess of non-target cells to immunologically remove phage antibodies directed against antigens present on non-target cells. This additional step may be performed either prior to incubating the phage antibody library with the cell population, or prior to isolating monoclonal antibodies from the enriched library.

Alternatively, it is possible to preincubate the cell population with a saturating amount of antibodies having known specificity prior to incubating the cell population with the phage antibody library. This immunologically blocks the known antigen binding sites on the target cell surface, and avoids enriching phage antibodies having the known specificity.

The goal of selectively enriching phage antibodies directed against antigens present only on target cells and not on non-target cells can also be achieved by DNA subtraction hybridization. In this method, two cell type specific phage antibody libraries are generated using the present inventive method, one for the target cells and one for the non-target cells. DNA encoding antibodies directed against antigens present on both cell types are then removed from the first library by hybridizing DNA from the first library to an excess of DNA from the second library and subsequently removing the annealed hybrids. The resulting

subtracted phage antibody library will then contain mostly antibodies directed against antigens unique to the target cells. Standard methods of DNA subtraction hybridization known to those skilled in the relevant art, or variations of these methods, may be used in this step.

5           The invention is further described in the following non-limiting examples.

## EXEMPLIFICATION

### Example 1

5 Laboratory mice are immunized against KG1a cells by injection of the cells intraperitoneally three times at intervals ranging from six weeks to six months. The KG1a cell line is an available human myeloblastoid cell line, and is characterized in several publications, including Koeffler et al., Blood 56, 265-273 (1980).

10 Six days following the last injection, spleen cells are isolated from the animals, lymphocytes are separated on a Ficoll-Paque™ gradient, commercially available from Pharmacia LKB Biotechnology, Inc., Piscataway, New Jersey, and total RNA is prepared using standard, known techniques. A combinatorial phage antibody library of  $10^7$  recombinants is prepared using standard protocols as described in Barbas et al., Methods 2, 119-124 (1991), and as described in further  
15 detail below.

KG1a cells are incubated with phage particles from the combinatorial library in binding buffer on ice for 30 minutes, and then spun through a calf serum cushion three times. Phage particles bound to the surface of the cells are eluted in an acid buffer (such as 0.1 M HCl, adjusted with glycine to pH 2.2; and  
20 BSA 1mg/ml), recovered, and used to infect XL1-Blue bacteria, available from Stratagene Cloning Systems, La Jolla, California. The phagemids in the infected bacteria are expanded in culture and rescued with helper phage VCSM13 to yield an enriched phage antibody library. The library is then used in another round of enrichment.

25 This enrichment process is repeated several times, four times in this example. Table 1 lists the number of phage particles recovered after each round of enrichment, showing a steady increase in the number of recovered phage particles after each round. In contrast, a negative control experiment (results not shown) using mouse thymocytes as the target cells for binding with the phage  
30 shows no increase in the number of phage particles recovered.

Table 1

Enrichment of phage antibodies by direct affinity selection on the cell surface

		<u>Round 1</u>	<u>Round 2</u>	<u>Round 3</u>	<u>Round 4</u>
5	No. phage recovered	$2 \times 10^5$	$8 \times 10^5$	$20 \times 10^5$	$20 \times 10^5$

Antigen binding fragments of the antibodies (Fab fragments) are produced from either the unenriched library or the final library after four rounds of enrichment and tested for reactivity to the KG1a cells by indirect fluorescence-activated cell sorting (FACS) analysis. FIGURES 3(a) through 3(d) are histograms showing fluorescence patterns of KG1a cells and mouse thymocytes stained with the unenriched (a,c) or enriched (b,d) antibody libraries. The abscissa indicates relative fluorescence intensity and the ordinate indicates relative cell number.

The staining of KG1a with the unenriched library (a) is no different from that of the background, whereas staining with the enriched library (b) gives signals approximately twenty times more intense. In contrast, the staining of mouse thymocytes with either the unenriched (c) or enriched (d) libraries are both indistinguishable from the background. The KG1a cells and mouse thymocytes have different autofluorescence patterns, thus accounting for the difference in background staining. These results demonstrate that phage antibody libraries can be enriched by the inventive method of direct affinity selection on the cell surface to generate monoclonal antibodies highly reactive to target cells.

To determine the fraction of monoclonal antibodies in the enriched phage antibody library that are specific for KG1a cells, Fab fragments are produced separately from ten individual clones picked at random from the library. FACS analysis of these samples (results not given) shows that the Fab fragments from all of the ten clones strongly react with KG1a cells and with variable patterns, indicating that all of the ten monoclonal antibodies are specific for KG1a and probably recognize at least several different surface antigens on the cells. By extension, the entire enriched antibody library most likely contains a large number of monoclonal antibodies recognizing many different surface antigens on KG1a cells.

### Example 2

KG1a cells, which bear the human CD34 antigen on their surface, are stained with CD34-specific antibodies according to Civin et al., J Immunol 133,

**THIS PAGE BLANK (USPTO)**

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the invention claimed herein.

5 All cited literature are incorporated by reference in their entirety.

We claim:

1. Method for generating monoclonal antibodies directed against previously uncharacterized and unpurified antigens on the surface of target cells in a cell population, using a combinatorial library of antibodies  
5 expressed on the surface of filamentous phage particles, the method comprising the steps of:
  - A. incubating the library with the target cells under conditions sufficient to bind a portion of the phage particles to the target cells;
  - B. separating the target cells and the bound phage particles from  
10 unbound phage particles;
  - C. recovering the phage particles bound to the target cells;
  - D. amplifying the bound phage particles to create an enriched library; and
  - E. isolating from the enriched library monoclonal antibodies  
15 specific for the target cells.
2. The method of claim 1, wherein the target cells comprise a phenotypically defined target cell subpopulation and the cell population comprises a heterogeneous cell population, the method further comprising,  
20 prior to Step A:
  - purifying the target cell subpopulation from the heterogeneous cell population.
3. The method of claim 1, wherein the target cells comprise a  
25 phenotypically defined target cell subpopulation and the cell population comprises a heterogeneous cell population, wherein Step B comprises:
  - copurifying the target cell subpopulation and the bound phage particles from the heterogeneous cell population and unbound phage particles.

4. The method of claim 3, wherein Step B further comprises:  
copurifying the target cells and the bound phage particles by one  
method selected from the group consisting of: fluorescence-activated cell  
5 sorting (FACS); immunological panning; magnetic bead separation; and,  
affinity chromatography.
5. The method of claim 1, wherein the target cells comprise a defined cell  
type in a frozen tissue section, and Step B comprises:  
10 identifying the target cells by immunohistochemistry; and  
microdissecting the target cells and the bound phage particles from the  
tissue section.
6. The method of claim 1, wherein the phage particles are phage  
15 antibodies, and wherein Step B comprises:  
identifying functionally important phage antibodies bound to the  
target cells by assaying for the consequences of a phage antibody-target cell  
binding using one assay selected from the group consisting of: measuring  
intracellular calcium concentration; detecting cell activation, cell proliferation,  
20 or cell killing; detecting blocking of cell activation, cell proliferation, or cell  
killing; and, analyzing expression of preselected genes.



## AMENDED CLAIMS

[received by the International Bureau on 24 October 1994 (24.10.94);  
original claims 6 and 7 replaced by new claim 6  
other claims unchanged (3 pages)]

1. Method for generating monoclonal antibodies directed against previously uncharacterized and unpurified antigens on the surface of target cells in a cell population, using a combinatorial library of antibodies expressed on the surface of filamentous phage particles, the method comprising the steps of:
  - A. incubating the library with the target cells under conditions sufficient to bind a portion of the phage particles to the target cells;
  - B. separating the target cells and the bound phage particles from unbound phage particles;
  - 10 C. recovering the phage particles bound to the target cells;
  - D. amplifying the bound phage particles to create an enriched library; and
  - E. isolating from the enriched library monoclonal antibodies specific for the target cells.
- 15 2. The method of claim 1, wherein the ~~target cells comprise a cell line~~,
3. The method of claim 1, wherein the target cells comprise a phenotypically defined target cell subpopulation and the cell population comprises a heterogeneous cell population, the method further comprising, prior to Step A:  
20 purifying the target cell subpopulation from the heterogeneous cell population.
4. The method of claim 1, wherein the target cells comprise a phenotypically defined target cell subpopulation and the cell population comprises a heterogeneous cell population, wherein Step B comprises:  
25 copurifying the target cell subpopulation and the bound phage particles from the heterogeneous cell population and unbound phage particles.
- 30 5. The method of claim 4, wherein Step B further comprises:

copurifying the target cells and the bound phage particles by one method selected from the group consisting of: fluorescence-activated cell sorting (FACS); immunological panning; magnetic bead separation; and, affinity chromatography.

- 5 6. The method of claim 1, wherein the target cells comprise a defined cell type in a frozen tissue section, and Step B comprises:

identifying the target cells by immunohistochemistry; and  
microdissecting the target cells and the bound phage particles from the  
tissue section.

10

7. The method of claim 1, wherein the phage particles are phage antibodies, and wherein Step B comprises:

identifying functionally important phage antibodies bound to the target  
cells by assaying for the consequences of a phage antibody-target cell binding  
15 using one assay selected from the group consisting of: measuring intracellular  
calcium concentration; detecting cell activation, cell proliferation, or cell killing;  
detecting blocking of cell activation, cell proliferation, or cell killing; and,  
analyzing expression of preselected genes.

AMENDED SHEET (ARTICLE 19)

8. The method of claim 1, further comprising the step of, prior to Step A:  
preadsorbing the phage antibody library with an excess of non-target cells,  
to immunologically remove phage antibodies directed against antigens present on  
non-target cells.

5

9. The method of claim 1, further comprising the step of, prior to Step A:  
preincubating the target cells with a saturating amount of preselected  
antibodies having known affinity in order to block binding sites on the target cells  
for the preselected antibodies.

10

10. The method of claim 1, further comprising the step of, prior to Step E:  
adsorbing the phage antibody library with an excess of non-target cells to  
immunologically remove phage antibodies directed against antigens present on  
non-target cells.

15

11. The method of claim 1, further comprising the step of, prior to Step E:  
removing, from the enriched library, phage antibodies directed against  
antigens present on non-target cells by DNA subtraction hybridization.

20



12. The method of claim 1, further comprising, the step of, after Step D,  
reiterating Step A through Step D to further enrich the library.



13. The cell-type specific phage antibody library produced by the process of  
claim 1.

25



14. The monoclonal antibodies generated from the cell type specific phage  
antibody library produced by the process of claim 1.

1 / 3

**FIG.\_1****FIG.\_2**



3 / 3



## INTERNATIONAL SEARCH REPORT

 tional application No.  
 PCT/US94/05124

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) :C07K 15/28

US CL :530/387.3, 867

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/387.3, 867

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS

SEARCH TERMS: PHAGE AND ANTIBODY

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim N .
Y	EP, A2, 0,460,607 (HELLSTROM ET AL.) 11 DECEMBER 1991, SEE ENTIRE DOCUMENT	1-6
Y	J. IMMUNOLOGY, VOL. 135, NO. 4, ISSUED OCTOBER 1985, LEDBETTER ET AL., "ANTIBODIES TO TP67 AND TP44 AUGMENT AND SUSTAIN PROLIFERATIVE RESPONSES OF ACTIVATED T CELLS," PAGES 2331-2336, SEE ENTIRE DOCUMENT.	1-6
Y	HARLOW ET AL., "ANTIBODIES A LABORATORY MANUAL" PUBLISHED 1988 BY COLD SPRING HARBOR LABORATORY (N.Y.), PAGES 496-497.	1-6

☒ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be of particular relevance	X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	A*	document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

30 JUNE 1994

Date of mailing of the international search report

29 JUL 1994

 Name and mailing address of the ISA/US  
 Commissioner of Patents and Trademarks  
 Box PCT  
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DONALD E. ADAMS PH.D.

Telephone No. (703) 308-0196

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US94/05124

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	NATURE, VOL. 308, ISSUED 08 MARCH 1984, HENDRICK ET AL., "ISOLATION OF CDNA CLONES ENCODING T CELL-SPECIFIC MEMBRANE-ASSOCIATED PROTEINS", PAGES 149-153, SEE ENTIRE DOCUMENT.	1-6